

Job Sheet for Seeding

Planned by:MO Mix 9 CP2

Date:

County:

Client:

Address:

Farm(s):

Tract(s):

Fields:

CIN#:

Acres:1.00

Program:

Primary / Cost-shared FOTG Practice:Wildlife Habitat Planting (420)

Secondary FOTG Practice:

Primary Purpose of Seeding:

Site Soil Group:

OR

Ecological Site:

Additional Information:

Seedbed Preparation for Grass and/or Forb/Legume Seeding

Current Conditions:

Conditions Desired at Planting:

☒ Row crop stubble or tilled seed bed (to incorporate amendments, level field, or following construction operations)

☐ Killed sod (renovation; for example from introduced cool season grasses to native warm season grasses or from warm season grasses to a pollinator forb only planting)

☐ Suppressed existing vegetation, as for an inter-seeding of legumes on pastureland or native forbs on natural communities

☐ Slopes that are 4:1 or steeper that will receive a critical area planting (broadcast planting method only)

Seed Mix Contains:

☐ Cool Season Grasses (majority)

☐ Warm Season Grasses (majority)

☐ Native Forbs

☐ Native/Introduced Legumes

Planting Method:

Interseeding:No

Planting Season:

Acceptable Planting Dates:

Optimal Planting Dates:

Seedbed Prep:

Seed Placement:

Additional Instructions:

Fertilizer and Lime

Planned

Is Fertilizer and/or Lime Needed?

☐ Program rules and/or planned practice(s) do not require soil amendments (example - general CRP plantings, wildlife plantings on cropland)

☐ Program rules and/or planned practices require soil amendments be applied according to a recent soil test (less than 4 years old and after the most recent application of liming material and/or fertilizer). Apply fertilizer and/or lime according to soil test recommendations as indicated below.

☐ Program rules and/or planned practices require soil amendments. Blanket rates will be used for a Critical Area Planting (342) as indicated below because a recent soil test is not available (as for constructed fills and finished grades).

Soil Test Recommendations

Salt pH from soil test results:

Lime*	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
lbs/ac			
Total Needed	Not req.	0	0

Blanket rates for constructed fills when a current soil test is not available:

lbs/ac	#N/A		
Total Needed	#N/A		

What legumes are included in the seed mix?

☐ None

☐ Alfalfa

☐ Cool Season Legumes other than Alfalfa

☐ Warm Season Legumes

When to apply lime:

When to apply fertilizer:

No fertilizer will be applied. Soil nutrients are within ranges required for the planned species that will be planted.

Page 1 of 6

Additional Information:

Companion Crop Specifications

A companion crop may be established prior to or at the same time as a permanent seeding of grass, forbs, and/or legumes.  
The companion crop will be selected and managed to minimize competition with the permanent seeding.

Planned Companion Crop:

☐ None Planned

☐ Redtop

☐ Spring Oats

☐ Winter Wheat

Rate per Acre (lbs):

Total Bulk Lbs to Seed:

Additional Information:

Applied

Companion Crop Planted:

☐ Redtop

☐ Spring Oats

☐ Winter Wheat

Bulk Lbs Planted:

Total Lbs	Lbs/Ac
	0.0

Seeding Mix Summary												
Acres to be Seeded:		1.00										
Practice Adjustment Factor:					Broadcast Adjustment Factor:			Interseeding Adjustment Factor: 1				
Apply Adjustment Factors to:												
Planned Seed Mix				Seed Mix Purchased and Planted								
Species	Acceptable Cultivar(s)	PLS lbs/ac	Total PLS lbs Needed	Bulk lbs Seeded	% Purity	% Germ	% Dormant/ Hard Seed	PLS lbs Planted	Shortage/ Excess	% of Mix	Cultivar OR Origin for MO Source Identified Class Seed	Meets all Require- ments
Little Bluestem, Schizachyrium scoparium	Aldous (statewide), Camper (north MO), Cimarron (south MO - sandy sites only), Ozark (statewide), Yellow Tag (statewide)	2.555	2.555						-2.555			
Switchgrass, Panicum virgatum	Alamo (south MO - lowland sites), Blackwell (statewide), Cave-In-Rock (statewide), Kanlow (statewide - lowland sites), Nebraska 28 (north MO), Pathfinder (north MO), Trailblazer (north MO), Yellow Tag (statewide)	1.675	1.675						-1.675			
Black-eyed Susan (B), Rudbeckia hirta		0.075	0.075						-0.075			
Canada Milkvetch, Astragalus canadensis		0.020	0.020						-0.020			
Common / Annual Lespedeza, Kummerowia stipulacea, K. striata		0.730	0.730						-0.730			
Common Evening Primrose (B), Oenothera biennis		0.030	0.030						-0.030			

False Sunflower, <i>Heliopsis helianthoides</i>		0.050	0.050						-0.050																						
Foxglove Beardtongue, <i>Penstemon digitalis</i>		0.019	0.019						-0.019																						
Golden Alexanders, <i>Zizia aurea</i>		0.070	0.070						-0.070																						
Gray Goldenrod, <i>Solidago nemoralis</i>		0.010	0.010						-0.010																						
Gray-headed Coneflower, <i>Ratibida pinnata</i>	Available: Sunglow (statewide)	0.055	0.055						-0.055																						
Illinois Bundleflower, <i>Desmanthus illinoensis</i>	Available: Reno Germplasm (statewide)	0.150	0.150						-0.150																						
Lanceleaf Coreopsis, <i>Coreopsis lanceolata</i>		0.100	0.100						-0.100																						
Maximilian Sunflower, <i>Helianthus maximiliani</i>		0.011	0.011						-0.011																						
Pale Purple Coneflower, <i>Echinacea pallida</i>		0.030	0.030						-0.030																						
Partridge Pea (A), <i>Chamaecrista fasciculata</i>	Available: Riley (statewide), Lark Selection (statewide)	0.100	0.100						-0.100																						
Prairie Cinquefoil, <i>Drymocallis arguta</i>		0.010	0.010						-0.010																						
Purple Coneflower, <i>Echinacea purpurea</i>		0.135	0.135						-0.135																						
Purple Prairie Clover, <i>Dalea purpurea</i>	Available: Kaneb (statewide)	0.127	0.127						-0.127																						
Rattlesnake Master, <i>Eryngium yuccifolium</i>		0.020	0.020						-0.020																						
Roundhead Lespedeza, <i>Lespedeza capitata</i>	Available: Kanoka (statewide)	0.020	0.020						-0.020																						
Showy Ticktrefoil, <i>Desmodium canadense</i>	Available: Alexander Germplasm (statewide)	0.040	0.040						-0.040																						
Smooth Aster, <i>Symphyotrichum laeve</i>		0.010	0.010						-0.010																						
Stiff Goldenrod, <i>Oligoneuron rigidum</i> , <i>Solidago rigida</i>		0.010	0.010						-0.010																						
Upright Prairie Coneflower, <i>Ratibida columnifera</i>		0.050	0.050						-0.050																						
White Prairie Clover, <i>Dalea candida</i>		0.023	0.023						-0.023																						
Wild Bergamot, <i>Monarda fistulosa</i>		0.048	0.048						-0.048																						
Total Planned:		6.173	6.173																												
Planned Grasses PLS/ft <sup>2</sup> :		24.08	Total PLS lbs. Forbs Planted: _____																												
Planned Forbs/Legumes PLS/ft <sup>2</sup> :		15.99	Total % of Mix: _____																												
Planned Total PLS/ft <sup>2</sup> :		40.07	% of mix of annual/biennial species: _____																												
Customer and Seed Dealer Please Note:																															

Only viable, high quality and adapted seed will be used. All seed shall have a current seed test within 10 months of the planting date that lists germination, purity, and hard seed as a percentage for determining pure live seed and lists the percent of weed seed present that meets State seed quality law standards. Seed test results must be provided to the customer and USDA for confirmation that seed planted meets specifications outlined in this plan. Seed must be clean and relatively free of weed seed and contaminants. Seed that has become wet, moldy, or otherwise damaged in transit or storage is not acceptable. Certified seed is preferred but not required unless Missouri Source Identified Class Seed, or yellow tag, is denoted as being required.

Legume seed shall be inoculated with the proper, viable Rhizobium bacteria species prior to planting according to Missouri Agronomy Technical Note MO-36, "Legume Inoculation." Pre-inoculated seed shall be planted prior to the expiration date on the inoculum tag or be re-inoculated with the appropriate inoculum within 24 hours prior to seeding. Inoculated seed will not be exposed to direct sunlight. Inoculation of legumes may be waived when a current pasture inventory (JS-AGRON-24 or similar inventory) shows that the legume species to be seeded occupies more than 5 percent of the vegetative cover on the site prior to seeding.

For "Acceptable Cultivar(s)," north Missouri (MO) is all counties north of Bates, Henry, Benton, Morgan, Moniteau, Cole, Osage, Gasconade, Franklin, and St. Louis Counties. South Missouri is all counties including and south of those listed. Refer to Native Cultivars or Selections Information Sheet (IS-MO723N) for guidance on selecting the appropriate native grass cultivars to meet objectives. In all cases where a cultivar(s) is/are noted under the Acceptable Cultivar(s) heading above, Missouri Source Identified Class Seed (yellow tag seed) may be used instead of a cultivar variety or selection.

**The following information applies only if "Locally adapted Missouri Source Identified Class Seed (yellow tag) required" is shown under the "Acceptable Cultivar(s)" heading above:** Locally adapted Missouri Source Identified Class Seed, or yellow tag, is defined as a native plant that source genetically originated in Missouri, was not introduced, and existed within the state borders prior to arrival of settlers. All seed from herbaceous material shall comply with Missouri seed laws including Missouri Crop Improvement Association guidance. All seed will comply with AOSCA (Association of Official Seed Certifying Agencies) certification procedures (including appropriate tagging) to include third-party verification by the Missouri Crop Improvement Association of source, genetic identity, and genetic purity of wildland collected or field or nursery grown plant germplasm materials. Seed must be Missouri origin (grown in Missouri) and certified as Missouri Source Identified Class. Source Identified Certification means: parent seed is collected from natural remnant Missouri populations; no selection, testing, or breeding for specific traits; production fields are inspected to verify species, source, and lack of noxious weeds; and seed is certified for purity and germination. If Missouri origin (grown) source identified class seed is not available, Missouri source identified class seed may be obtained from adjoining states.

<b>Mulching Specifications (After Grass Seeding)</b>			
Type of Mulch Required: <input type="checkbox"/> None <input type="checkbox"/> Straw, Grass Hay, Alfalfa Hay - 1 1/2 to 2 1/2 Tons/Acre			
Amount of Mulch Required for This Site:			
Minimum Rate (lbs):	3,000	OR	Tons: 1.50
Maximum Rate (lbs):	5,000	OR	Tons: 2.5
Additional Information: n/a			
<b>Applied</b>			
Type of Mulch Used:			
Number of Bales:		Bale Weight, lbs:	
Mulch Applied, Tons:		Excess/Shortage, Tons:	
Based on the information in this job sheet, all requirements for this seeding should be completed by: _____ Year: _____			
Was site prep completed as outlined above? _____			
Seeding was completed on _____ according to the above requirements.			
(Date)			
Producer's or Contractor's Signature		Date	
<b>For NRCS Use Only:</b>			
Does this seeding meet all NRCS standards and specifications and was implemented as planned above? _____			
Certified by: _____			
NRCS Representative		Date	Field Office
Additional Comments: _____			
When seeding is completed, return seeding plan to the Natural Resources Conservation Service with all receipts pertaining to this seeding (seed, fertilizer, lime, mulch, companion crop, labor, drill rental, etc.) and seed tests, including tags for yellow tag seed.			